

Enhancing Coffee Quality Through Modern Technology:A Focus on Ultrasound

Sara Malakouti^{1*}, Ahad Rahbari Maleki², Milad Shaghaghy Ranjbar³

1- Lecturer of Shabestar University of Applied Sciences

2- student of food science and technology, Shabestar University of Applied Sciences University, Tabriz

3- Lecturer of Shirin Asal University of Applied Sciences

Abstract

Coffee, a globally significant commodity, undergoes a meticulous journey from cultivation to the final brewed cup. This article explores the critical processes of planting, processing, roasting, and their impact on coffee quality. Optimal cultivation practices ensure high yields and premium cherries, while effective harvesting and post-harvest handling preserve coffee integrity. Processing methods, such as wet, semi-dry, and dry, significantly influence flavor profiles, affecting attributes like acidity, body, and aroma. Roasting, a pivotal step, develops distinct flavors and aromas requiring precise control. Integrating ultrasound technology in coffee processing enhances quality through cleaning, preventing biofilm formation, and optimizing extraction efficiency.

Keywords: Coffee cultivation, coffee processing, roasting, coffee quality, Arabica, Robusta, flavor profile, ultrasound, food processing, biofilm control.

Introduction

Coffee stands as a cornerstone of global agricultural and economic landscapes, revered not only for its stimulating caffeine content but also for its rich array of bioactive compounds and antioxidant properties. The journey of coffee from its cultivation to the final cup undergoes a series of intricate processes—planting, processing, storing, roasting, and grinding—that profoundly influence its complex aroma and flavor profile. These processes are not merely functional but crucial in shaping coffee's sensory characteristics, making them subjects of meticulous attention and innovation in modern coffee production.

Advancements in technology, such as ultrasound, play a pivotal role in enhancing the efficiency and quality of food processing, particularly in coffee production. Ultrasound technology's application extends from improving cleaning processes to preventing biofilm formation in food processing equipment, thereby safeguarding the integrity and hygiene standards crucial to maintaining the quality of coffee products.

Background

Coffee, from the genus *Coffea* in the Rubiaceae family, encompasses over 100 species, though only *Coffea arabica* (Arabica) and *Coffea canephora* (Robusta) are commercially significant. Arabica, known for its sweet, floral/fruity taste, dominates global production, constituting about 70% of coffee production worldwide.

Coffee Production and Consumption

Second only to oil exports, coffee production thrives in tropical regions with consumption largely in developed countries. This distribution owes to favorable climates and economic factors that facilitate processing and re-exporting in northern nations.

From Farm to Cup: Factors Affecting Coffee Quality

Coffee bean quality hinges on environmental factors like climate, influencing acidity, texture, sweetness, and aroma. Variations in flowering times due to water stress and temperature changes produce both ripe and unripe fruits at harvest, impacting overall bean quality.

Coffee Fruit Processing

Essential for transportation, coffee fruit processing discards 80-90% of fruit, primarily pulp and skin. Processing methods—wet, semi-dry, and dry—greatly influence coffee flavor profiles, modifying attributes like acidity, body, and aroma.

- **Wet Method:** Ideal for specialty coffees, this costly process uses ripe, whole fruits, yielding aromatic, less bitter coffee. However, environmental concerns arise from wastewater high in organic material and acidity.
- **Semi-Dry or Pulped Natural Method:** This method reduces acidity while enhancing body, ideal for aroma and viscosity, with lower water consumption, promoting environmental sustainability.
- **Dry or Natural Method:** The most widespread method globally, it involves air-drying coffee fruits over four weeks, potentially reducing aroma quality due to volatile compound loss.

Coffee Roasting

Critical for creating specific sensory characteristics, roasting involves mass transfer via heating and subsequent endothermic and exothermic reactions. Proper storage ensures green and roasted bean quality, crucial for retaining freshness and flavor.

Roasting Processes and Control

Roasting methods vary across domestic, commercial, and industrial scales, employing indirect (convection) or direct (conduction) heating. Varying standards globally influence coffee flavor and aroma profiles significantly.

Complex Aroma of Coffee

The Maillard reaction and caramelization are pivotal in developing coffee's rich aroma and flavor. These reactions, combining sugars and amino acids, produce diverse flavor compounds contributing to coffee's complex profile.

Ultrasound in Food Processing

Ultrasound, particularly low-frequency, high-energy types, enhances food processing efficiency and quality. Its applications range from drying, freezing, and thawing processes to cleaning and biofilm prevention in food processing environments.

Ultrasound Cleaning and Extraction

Known for effective cleaning via microjet formation during cavitation, ultrasound enhances bioactive compound extraction from food materials. It efficiently removes surface lipids and enhances delicate food cleaning processes.

Biofilm Control with Ultrasound

Biofilms in food processing equipment pose challenges to food safety and quality. Ultrasound, a nonchemical, environmentally friendly technology, effectively prevents and reduces biofilm formation, crucial for maintaining food processing hygiene.

Airborne Ultrasound and Food Processing

Less common but effective, airborne ultrasound aids drying and particulate removal, enhancing air quality in processing environments. It accelerates drying without compromising product quality, as observed in apple slices and vegetable drying processes.

Effect of Ultrasound on Freezing Rates

Ultrasound accelerates ice crystal formation, yielding smaller crystals that preserve food quality. It enhances nucleation and heat transfer, minimizing cellular structure damage during freezing, crucial for maintaining food texture.

Defrosting and Thawing with Ultrasound

Ultrasound-assisted thawing accelerates thawing processes, reducing drip loss and enhancing food quality. Acoustic energy absorption by frozen foods at specific frequencies improves thawing rates, crucial for preserving food texture and quality.

Viscosity Reduction with Ultrasound

Controlling product viscosity during production improves processing efficiency, reducing fouling and contamination risks. Ultrasound-induced cavitation shear forces can temporarily alter viscosity, benefiting processes like spray drying and homogenization.

Dehydration and Drying with Ultrasound

Critical for food preservation, ultrasound enhances drying rates and efficiency while maintaining product quality. It improves conventional drying methods' limitations, preserving heat-sensitive materials effectively.

Ultrasound and Osmotic Dehydration

Ultrasound creates microscopic channels in fruit structures, enhancing osmotic dehydration efficiency without compromising fruit sugars. Combined with osmotic methods, ultrasound accelerates dehydration rates while preserving fruit quality.

Ultrasound Extraction of Bioactive Compounds

Ultrasound's effectiveness at lower temperatures makes it ideal for extracting bioactive compounds prone to degradation. Factors like extraction time, temperature, and ultrasonic intensity affect yield, benefiting extraction from fruits, seeds, and plants.

Conclusion

The meticulous journey of coffee from plant to cup underscores the importance of each step in the process. From selecting the right variety and cultivating under optimal conditions to carefully harvesting and processing the cherries, each phase plays a critical role in defining the final quality of the coffee. Proper cultivation practices ensure healthy plant growth, optimal yield, and high-quality cherries. Effective harvesting and post-harvest handling maintain the integrity and quality of the coffee cherries, preventing spoilage and preserving the desirable flavor characteristics. Understanding and optimizing each step of the coffee production process can lead to the production of superior coffee, appreciated for its rich aroma and complex flavor. Additionally, integrating advanced technologies like ultrasound in food processing enhances product quality and processing efficiency, supporting the sustainability of the coffee and food industries. Ultrasound technology's role in preventing biofilm formation and enhancing various food processing stages highlights its potential for broader application in improving food safety and quality.

References

- Achat, S., Tomao, V., Madani, K., Chibane, M., Elmaataoui, M., Dangles, O., and Chemat, F. 2012. Direct enrichment of olive oil in oleuropein by ultrasound-assisted maceration at laboratory and pilot plant scale. *Ultrason. Sonochem.* 19(4):777–786.
- Ahmad, A. L., Che Lah, N. F., Ismail, S., and Ooi, B. S. 2012. Membrane antifouling methods and alternatives: Ultrasound approach. *Sep. Purif. Rev.* 41(4):318–346.
- Ahmad-Qasem, M. H., Cánovas, J., Barrajón-Catalán, E., Micol, V., Cárcel, J. A., and García-Pérez, J. V. 2013. Kinetic and compositional study of phenolic extraction from olive leaves (var. Serrana) by using power ultrasound. *Innov. Food Sci. Emerg.* 17:120–129.
- Allaf, T., Tomao, V., Ruiz, K., and Chemat, F. 2013. Instant controlled pressure drop technology and ultrasound assisted extraction for sequential extraction of essential oil and antioxidants. *Ultrason. Sonochem.* 20(1):239–246.
- Allen, V. M., Whyte, R. T., Burton, C. H., Harris, J. A., Lovell, R. D., Atterbury, R. J., and Tinker, D. B. 2008. Effect of ultrasonic treatment during cleaning on the microbiological condition of poultry transport crates. *Brit. Poult. Sci.* 49(4):423–428.
- Assami, K., Pingret, D., Chemat, S., Meklati, B. Y., and Chemat, F. 2012. Ultrasound induced intensification and selective extraction of essential oil from *Carum carvi* L. seeds. *Chem. Eng. Process: Process Intensif.* 62:99–105.
- Aversa, M., Van der Voort, A. J., de Heij, W., Tournois, B., and Curcio, S. 2011. An experimental analysis of acoustic drying of carrots: Evaluation of heat transfer coefficients in different drying conditions. *Dry. Technol.* 29(2):239–244.
- Awad, T. S., Moharram, H. A., Shaltout, O. E., Asker, D., and Youssef, M. M. 2012. Applications of ultrasound in analysis, processing and quality control of food: A review. *Food. Res. Int.* 48(2):410–427.
- Bantle, M., and Eikevik T. M. 2011. Parametric study of high-intensity ultrasound in the atmospheric freeze drying of peas. *Dry. Technol.* 29(10):1230–1239.
- Benito-Román, Ó., Alonso E., and Cocero M. J. 2013. Ultrasound-assisted extraction of β -glucans from barley. *LWT—Food Sci. Technol.* 50(1):57–63.
- Bhaskaracharya, R. K., Kentish, S., and Ashokkumar, M. 2009. Selected applications of ultrasonics in food processing. *Food Eng. Rev.* 1(1):31–49.
- Bimakr, M., Rahman, R. A., Taip, F. S., Adzahan, N. M., Islam Sarker, M. Z., and Ganjloo, A. 2012. Optimization of ultrasound-assisted extraction of crude oil from winter melon (*Benincasa hispida*) seed using response surface methodology and evaluation of its antioxidant activity, total phenolic content and fatty acid composition. *Molecules.* 17(10):11748–11762.
- Cacciola, V., Batllò, I. F., Ferraretto, P., Vincenzi, S., and Celotti, E. 2013. Study of the ultrasound effects on yeast lees lysis in winemaking. *Eur. Food Res. Technol.* 236(2):311–317.
- Chandrapala, J., Oliver, C., Kentish, S., and Ashokkumar, M. 2012. Ultrasonics in food processing—food quality assurance and food safety. *Trends Food Sci. Technol.* 26(2):88–98.
- Chandrapala, J., Oliver, C. M., Kentish, S., and Ashokkumar, M. 2013. Use of power ultrasound to improve extraction and modify phase transitions in food processing. *Food Rev. Int.* 29(1):67–91.
- Chemat, F., and Khan, M. K. 2011. Applications of ultrasound in food technology: Processing, preservation and extraction. *Ultrason. Sonochem.* 18(4):813–835.
- Chemat, S., and Esveld, E. D. 2013. Contribution of microwaves or ultrasonics on carvone and limonene recovery from dill fruits (*Anethum graveolens* L.). *Innov. Food Sci. Emerg.* 17:114–119.
- Clodoveo, M. L., Durante, V., and La Notte, D. 2013. Working towards the development of innovative ultrasound equipment for the extraction of virgin olive oil. *Ultrason. Sonochem.* 20(5):1261–1270.
- Delgado, A. E., Zheng, L., and Sun, D. W. 2009. Influence of ultrasound on freezing rate of immersion-frozen apples. *Food Bioprocess. Technol.* 2(3):263–270.
- Fahmi, R., Khodaiyan, F., Pourahmad, R., and Emam-Djomeh, Z. 2011. Effect of ultrasound assisted extraction upon the protein content and rheological properties of the resultant soymilk. *Adv. J. Food Sci. Technol.* 3(4):245–249.
- Fahmi, R., Khodaiyan, F., Pourahmad, R., and Emam-Djomeh, Z. 2014. Effect of ultrasound assisted extraction upon the Genistin and Daidzin contents of resultant soymilk. *J. Food Sci. Technol.* 51(10):2857–2861.



22. Fernandes, F. A., and Rodrigues, S. 2012. Ultrasound as pre-treatment for drying of genipap (*Genipa americana* L.). *Int. J. Food Eng.* 8(3):1556–3758.
23. Fernandes, F. A., Gallão, M. I., and Rodrigues, S. 2009. Effect of osmosis and ultrasound on pineapple cell tissue structure during dehydration. *J. Food Eng.* 90(2):186–190.
24. Fernandes, F. A., Rodrigues, S., Law, C. L., and Mujumdar, A. S. 2011. Drying of exotic tropical fruits: A comprehensive review. *Food Bioprocess. Technol.* 4(2):163–185.
25. Frias, J., Peñas, E., Ullate, M., and Vidal-Valverde, C. 2010. Influence of drying by convective air dryer or power ultrasound on the vitamin C and β -carotene content of carrots. *J. Agric. Food Chem.* 58(19):10539–10544.
26. Fu, Q., Zhang, L., Cheng, N., Jia, M., and Zhang, Y. 2014. Extraction optimization of oleanolic and ursolic acids from pomegranate (*Punica granatum* L.) flowers. *Food Bioprod. Process.* 92(3):321–327.
27. Gaete-Garretón, L., Vargas-Hernández, Y., Cares-Pacheco, M. G., Sainz, J., and Alarcón, J. 2011. Ultrasonically enhanced extraction of bioactive principles from Quillaja saponaria Molina. *Ultrasonics.* 51(5):581–585.
28. Gallego-Juárez, J. A., Riera, E., De la Fuente Blanco, S., Rodríguez-Corral, G., Acosta-Aparicio, V. M., and Blanco, A. 2007. Application of high-power ultrasound for dehydration of vegetables: Processes and devices. *Dry. Technol.* 25(11):1893–1901.
29. Garcia-Noguera, J., Oliveira, F. I., Weller, C. L., Rodrigues, S., and Fernandes, F. A. 2014. Effect of ultrasonic and osmotic dehydration pre-treatments on the colour of freeze dried strawberries. *J. Food Sci. Technol.* 51(9):2222–2227.
30. García-Pérez, J. V., Cárcel, J. A., Riera, E., and Mulet, A. 2009. Influence of the applied acoustic energy on the drying of carrots and lemon peel. *Dry. Technol.* 27(2):281–287.
31. Garcia-Perez, J. V., Carcel, J. A., Riera, E., Rosselló, C., and Mulet, A. 2012. Intensification of low-temperature drying by using ultrasound. *Dry. Technol.* 30(11–12):1199–1208.
32. Garofulic, I. E., Jambrak, A. R., Milicevic, B., Dragovic-Uzelac, V., and Zoric, Z. 2012. The effect of high power ultrasound and gamma irradiation on phenolic compounds in pomegranate juice. *Ultrason. Sonochem.* 19(3): 634–641.
33. Gawron, P., Pellicer, N., Portero, P., Belda, M., and Graell, J. 2008. The use of ultrasound as a technology to control the fouling rate in crossflow filtration by modifying yeast cell morphology. *Desalination.* 228(1–3):127–135.
34. Grønnevik, H., Liaaen-Jensen, S., and Aksnes, A. 2011. Effect of ultrasound on quality of fish oil. *Eur. J. Lipid Sci. Technol.* 113(5):537–546.
35. Ignat, A., Manzocco, L., Brunton, N., Nicoli, M. C., and Lyng, J. G. 2014. The effect of pulsed electric field pre-treatment on enrichment of antioxidant activity extractability from yellow onions. *Innov. Food Sci. Emerg.* 21: 58–63.
36. Jiménez, A., Beltrán, G., and Uceda, M. 2007. High-power ultrasound in olive paste pretreatment: Effect on process yield and virgin olive oil characteristics. *Ultrason. Sonochem.* 14(6):725–731.
37. Kadkhodae, R., and Povey, M. J. 2008. Ultrasonic inactivation of *Bacillus* α -amylase. I. Effect of gas content and emitting face of probe. *Ultrason. Sonochem.* 15(2):133–142.
38. Kadkhodae, R., and Povey, M. J. 2008. Ultrasonic inactivation of *Bacillus* α -amylase. II. Effect of temperature. *Ultrason. Sonochem.* 15(2):143–151.
39. Karabacak, N., and Bozkurt, H. 2008. Effect of ultrasound and temperature on the rheological and physical properties of ice cream mixes. *J. Food Sci. Technol.* 45(2):194–199.
40. Kentish, S., Wooster, T. J., Ashokkumar, M., Balachandran, S., Mawson, R., and Simons, L. 2008. The use of ultrasonics for nanoemulsion preparation. *Innov. Food Sci. Emerg.* 9(2):170–175.
41. Lee, H., and Feng, H. 2011. Effects of ultrasound and pressure on enzymatic activity of pectinmethylesterase extracted from Valencia orange. *Innov. Food Sci. Emerg.* 12(4):467–471.
42. Lee, H., Zhou, B., Liang, X., and Feng, H. 2009. Inactivation of *E. coli* K12 in apple cider by combination of pulsed electric fields and ultrasound. *Food Bioproc. Technol.* 2(4):364–374.
43. Ma, Y., Ye, X., Hao, Y., Xu, G., Xu, G., and Liu, D. 2008. Ultrasound-assisted extraction of hesperidin from Penggan (*Citrus reticulata*) peel. *Ultrason. Sonochem.* 15(3):227–232.
44. McClements, D. J. 2011. Ultrasonic characterization of foods and drinks: Principles, methods, and applications. *Crit. Rev. Food Sci. Nutr.* 41(5):419–424.
45. Medina-Torres, L., Ayora-Talavera, T., Espinosa-Andrews, H., Sánchez-Contreras, A., and Pacheco, N. 2017. Ultrasound assisted extraction for the recovery of phenolic compounds from vegetable sources. *Agriculture.* 7(10):70.